

Patient: _____ Date of exam: _____

Central incisor position:

- Acceptable and to be reproduced in the definitive restoration.
- Needs correction at or before the definitive prosthesis(es).
- Not applicable/unable to assess.

Central display at rest: _____ mm (average male: 0-2 mm, average female 2-4 mm).

High smile line:

- High (>100% incisal display) (~10% of population, twice as common in females).
- Moderate (75-100% incisal display) (~70% of population).
- Low (<75% incisal display) (~20% of patients).
- Not applicable/unable to assess.

Vertical dimension of occlusion:

- maintained and adequate.
- seemingly maintained but inadequate.
- lost.

Occlusal plane:

- Acceptable
- Complex:

Function/parafunction/force concern:

- High (low FMA, deep bite, recognized parafunction, wear facets)
- Average (FMA is WNL, no known parafunction)
- LOW (High FMA, opposing denture, no known parafunction, history of long-standing long-span FDPs)

Support:

- Facial support necessary or likely necessary
- Lip support necessary or likely necessary

Clinical records checklist

- Clinical images (6 images)
- Intraoral scan
- Facebow transfer
- Pantomograph
- CBCT

Determinations

- Surgical consultation necessary to further prosthetic discussion.
- Mounted study models necessary to finalize treatment plan.